

DTIC FILE COPY

4

OFFICE OF NAVAL RESEARCH

Grant N00014-90-J-1193

TECHNICAL REPORT No. 6

AD-A219 436

Optical Phonons and Electronic Properties in Double Heterostructures

by

D. L. Lin, R. Chen and Thomas F. George

Prepared for Publication

in

Quantum Well and Superlattice Physics III

Edited by G. H. Döhler, E. Koteles and J. N. Schulman

Proceedings of the Society of Photo-Optical Instrumentation Engineers

Departments of Chemistry and Physics  
State University of New York at Buffalo  
Buffalo, New York 14260

March 1990

Reproduction in whole or in part is permitted for any purpose of the  
United States Government.

This document has been approved for public release and sale;  
its distribution is unlimited.

DTIC  
ELECTE  
MAR 16 1990  
S E D

00 03 15 034

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

## REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFFALO/DC/90/TR-6			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York		6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State, and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260			7b. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Grant N00014-90-J-1193		
8c. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.
					WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Optical Phonons and Electronic Properties in Double Heterostructures					
12. PERSONAL AUTHOR(S) D. L. Lin, R. Chen and Thomas F. George					
13a. TYPE OF REPORT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) March 1990	
				15. PAGE COUNT 15	
16. SUPPLEMENTARY NOTATION Prepared for publication in Quantum Well and Superlattice Physics III Edited by G. H. Döhler, E. S. Koteles and J. N. Schulman, Proceedings of the Society of Photo-Optical Instrumentation Engineers					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	OPTICAL PHONONS DOUBLE HETEROSTRUCTURES		
			ELECTRONIC PROPERTIES CONTINUUM MODEL		
			SEMICONDUCTOR INTERFACE MODES		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Optical phonon modes in a semiconductor double heterostructure (DHS) are examined within the continuum model. The interface modes found here can account for the novel phenomena observed in right-angle Raman scattering. The Hamiltonian for the electron interaction with phonon eigenmodes is derived and employed to study polaronic effects in the DHS. The $1s-2p_z$ transition energy of a magnetopolaron bound a hydrogenic impurity in the quantum well is calculated, and excellent agreement with experiments is obtained.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson			22b. TELEPHONE (Include Area Code) (202) 696-4410		22c. OFFICE SYMBOL

DD Form 1473, JUN 86

Previous editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE

UNCLASSIFIED

## Optical phonons and electronic properties in double heterostructures

D. L. Lin, R. Chen and Thomas F. George

Department of Physics and Astronomy  
State University of New York at Buffalo  
Buffalo, New York 14260

### ABSTRACT

Optical phonon modes in a semiconductor double heterostructure (DHS) are examined within the continuum model. The interface modes found here can account for the novel phenomena observed in right-angle Raman scattering. The Hamiltonian for the electron interaction with phonon eigenmodes is derived and employed to study polaronic effects in the DHS. The  $1s-2p_+$  transition energy of a magnetopolaron bound to a hydrogenic impurity in the quantum well is calculated, and excellent agreement with experiments is obtained.

### 1. INTRODUCTION

It is by now well known that in polar crystals such as III-V compounds, the electron-optical-phonon interaction plays an important role in determining the electronic properties at higher temperatures. Since these compounds are the usual materials used in the fabrication of microstructures such as heterostructures and superlattices, there has been growing interest in the study of electron-phonon interactions in these structures. Until recently, the bulk Fröhlich Hamiltonian has been employed in almost all calculations, even though the presence of surfaces and interfaces is expected to change the nature of lattice vibration modes dramatically.<sup>1,2</sup>


Although surface optical (SO) phonon modes have been discussed some time ago,<sup>3,4</sup> they have not been included until fairly recently in the treatment of polarons confined in a slab<sup>5</sup> or near the interface in semi-infinite systems.<sup>6,7</sup> In recent papers,<sup>8,9</sup> the optical phonon modes in a double heterostructure (DHS) of polar crystals have been solved within the continuum model. It is found that there are only two types of phonon modes: the interface (IN) and the confined bulk modes. The latter may be either longitudinal (LO) or transversal (TO) phonon modes. Therefore, the Fröhlich Hamiltonian involving only the bulk LO phonon cannot be applied to quantum wells and other microstructures.

We report in this paper the Hamiltonian derived for the electron-optical-phonon interaction in a DHS. Both the interface and bulk LO phonons are included. The interaction Hamiltonian is then employed to study the phonon effects on electronic and optical properties in the DHS. The self-energy of a free polaron is calculated for arbitrary well width. It approaches the bulk value of the side material in the limit of  $d = 0$ , and that of the central layer in the limit of  $d \rightarrow \infty$ . For a magnetopolaron bound to a hydrogenic impurity in the GaAs/AlGaAs quantum

well, the transition energy calculated from this Hamiltonian exhibits the strange pinning phenomena in strong magnetic fields as observed experimentally.

## 2. OPTICAL PHONON MODES

The detailed solution of phonon eigenmodes can be found in Refs. 8 and 9. Here we just outline what is essential for our discussion of polaronic effects. The geometry of our DHS is shown in Fig. 1. Material  $\nu$  ( $\nu = 1, 2$ ) is characterized



Accession For	
1. IS GRA&I	<input checked="" type="checkbox"/>
DMC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

Figure 1. Geometry of the double heterostructure.

by the dielectric constants  $\epsilon_\nu(\omega)$ ,  $\epsilon_{0\nu}$  and  $\epsilon_{\infty}$ , plasma frequency  $\omega_{p\nu}$ , number of ion pairs  $n_\nu$  per unit cell, natural frequency  $\omega_{0\nu}$ , and polarizability  $\alpha_\nu$ . The polarization vector  $\vec{P}$  is related to the relative displacement  $\vec{u}(\vec{r})$  of the ion pair and to the macroscopic field  $\vec{E}$ :

$$\begin{aligned}\vec{P}(\vec{r}) &= n_\nu e^* \eta_\nu^{-1/2} \vec{u}(\vec{r}) \\ &= \frac{1}{4\pi} \chi_\nu(\omega) \vec{E},\end{aligned}\quad (1)$$

where we have defined  $\eta_\nu^{1/2} = [1 + \alpha_\nu n_\nu (\lambda_{0\nu} - \lambda_\nu)]^{-1}$  and

$$\lambda_{0\nu} = 4\pi\omega_{0\nu}^2/\omega_{p\nu}^2, \quad \lambda_\nu = 4\pi\omega^2/\omega_{p\nu}^2. \quad (2a,b)$$

The dielectric susceptibility  $\chi_\nu(\omega)$  is given by

$$4\pi\chi_\nu^{-1} = (\lambda_{0\nu} - \lambda_\nu)\eta_\nu^2 - 4\pi/3 \quad (3)$$

and is related to the dielectric function by  $\chi_\nu(\omega) = \epsilon_\nu(\omega) - 1$ . With the frequency parameters (2), we find from (3)

$$\epsilon_\nu(\omega) = \epsilon_{\infty\nu}(\omega_{L\nu}^2 - \omega^2)/(\omega_{T\nu}^2 - \omega^2) \quad (4a)$$

$$\epsilon_{\infty\nu} = 1 + 4\pi n_\nu \alpha_\nu / (1 - \frac{4\pi}{3} n_\nu \alpha_\nu) \quad (4b)$$

The electrostatic potential  $\phi(\vec{r})$  that gives rise to the field  $\vec{E}$  satisfies

$$\vec{E}(\vec{r}) = -\nabla\phi(\vec{r}) \quad (5)$$

On the assumption that the dielectric function  $\epsilon_\nu(\omega)$  does not depend upon the spatial coordinates within each layer, we have from (5) and  $\nabla \cdot \vec{D} = 0$

$$\epsilon_\nu(\omega) \nabla^2 \phi(\vec{r}) = 0 \quad (6)$$

Since the DHS does not possess translational invariance in the z-direction, the two-dimensional (2D) vectors  $\vec{\kappa}$  and  $\vec{\rho}$  are introduced in the wave vector  $\vec{k} = (\vec{\kappa}, q)$  and in the position vector  $\vec{r} = (\vec{\rho}, z)$ . The potential can then be written as

$$\phi(\vec{r}) = \int d\vec{\kappa} \phi(\vec{\kappa}, z) e^{i\vec{\kappa} \cdot \vec{\rho}} \quad (7)$$

Substituting (7) in (6), we find that the Fourier components of the potential satisfy the equation

$$\epsilon_\nu(\omega) \left[ \frac{\partial^2}{\partial z^2} - \kappa^2 \right] \phi(\vec{\kappa}, z) = 0 \quad (8)$$

which may be solved by assuming either

$$\epsilon_\nu(\omega) = 0 \quad , \quad \nu = 1, 2 \quad (9a)$$

or

$$\left[ \frac{\partial^2}{\partial z^2} - \kappa^2 \right] \phi(\vec{\kappa}, z) = 0 \quad (9b)$$

It is seen from (4a) that (9a) is satisfied by the LO frequencies  $\omega^2 = \omega_{L\nu}^2$ . Because  $\omega_{L1} \neq \omega_{L2}$ , the vibrations on both sides of the interface have different time dependences. The boundary conditions that both  $\phi$  and the electric displacement  $D_z$  must be continuous across the interfaces then require the confinement of the LO modes. On the other hand, Eq. (9b) yields the interface modes of vibration. The polarization eigenvectors of all these modes as well as the dispersion relations have been worked out in Ref. 8. It is also shown there that the novel slab modes observed in right-angle Raman scattering<sup>10</sup> can be interpreted as the interface modes. As discussed in Ref. 8, in the vicinity of the center of the first Brillouin zone, the antisymmetric interface modes, which are predominantly transversal, oscillate at longitudinal frequency while the symmetric interface modes are longitudinal but oscillate at transversal frequency. Using Eqs. (8) and (9), we find after a straightforward calculation electrostatic potentials corresponding to these modes. Thus for the confined LO modes, the Fourier components of the potential are

$$\phi_m(\vec{\kappa}, z) = 4\pi C_m(\vec{\kappa}) \begin{cases} \sin(m\pi z/d) & , \text{ for even } m \\ \cos(m\pi z/d) & , \text{ for odd } m \end{cases} \quad (10a)$$

with the normalization constant

$$|C_m(\kappa)| = \omega_{p\nu} \eta_\nu^{-1/2} \sqrt{2/d} [\kappa^2 + (m\pi/d)^2]^{-1/2} \quad (10b)$$

Corresponding to the interface modes, the potentials are

$$\phi_{sj}(\vec{\kappa}, z) = \begin{cases} C_{sj} e^{-\kappa|z|} & , \quad |z| > d/2 \\ C_{sj} (1 + \epsilon_2/\epsilon_1) \cosh(\kappa z) & , \quad |z| < d/2 \end{cases} \quad (11a)$$

for the symmetric modes and

$$\phi_{aj}(\vec{\kappa}, z) = \begin{cases} -C_{aj} \text{sgn}(z) e^{-\kappa|z|} & , \quad |z| > d/2 \\ C_{aj} (1 + \epsilon_2/\epsilon_1) \sinh(\kappa z) & , \quad |z| < d/2 \end{cases} \quad (11b)$$

for the antisymmetric modes. The function  $\text{sgn}(z) = 1$  for  $z > 0$  and  $-1$  for  $z < 0$ . The index  $j = 1, 2$  in Eq. (11) labels the two branches of the symmetric or antisymmetric interface modes which have the same normalization constant as discussed in Ref. 8. The normalization constants are given by

$$|C_{sj}(\vec{\kappa})| = \frac{4\pi}{\sqrt{A}} \epsilon \kappa^{d/2} / \left\{ 2\kappa \left[ \frac{\eta_1 \chi_1^2}{\omega_{p1}^2} \tan\left(\frac{\kappa d}{2}\right) + \frac{\eta_2 \chi_2^2}{\omega_{p2}^2} \right] \right\}^{1/2} \quad (12a)$$

$$|C_{aj}(\vec{\kappa})| = \frac{4\pi}{\sqrt{A}} e^{\kappa d/2} \sqrt{\tanh(\frac{\kappa d}{2})} / \left\{ 2\kappa \left[ \frac{\eta_1 \chi_1^2}{\omega_{p1}^2} + \frac{\eta_2 \chi_2^2}{\omega_{p2}^2} \tanh(\frac{\kappa d}{2}) \right] \right\}^{1/2} \quad (12b)$$

### 3. ELECTRON-PHONON INTERACTIONS

To derive the Hamiltonian operator, we start with the free polarization or the free phonon. In terms of the polarization vector  $\vec{\pi}$  and its canonical conjugate

$\dot{\vec{\pi}}$ , we can write the Hamiltonian in the form

$$H_{ph} = \frac{1}{2} \sum_{\alpha} \int d\vec{r} \frac{4\pi\eta_{\nu}}{\omega_{p\nu}^2} [\dot{\vec{\pi}}_{\alpha}^*(\vec{r}) \dot{\vec{\pi}}_{\alpha}(\vec{r}) + \omega_{\alpha}^2 \vec{\pi}_{\alpha}^*(\vec{r}) \vec{\pi}_{\alpha}(\vec{r})] \quad (13)$$

$\vec{\pi}$  and  $\dot{\vec{\pi}}$  can be regarded as quantum mechanical operators and can be expressed in terms of their Fourier components as

$$\begin{aligned} \vec{\pi}(\vec{r}) = & \sum_{\vec{\kappa}, m} \sqrt{\hbar/8\pi A \omega_{L\nu}} [a_m(\vec{\kappa}) + a_m^{\dagger}(-\vec{\kappa})] \vec{\pi}_m(\vec{r}) \\ & + \sum_{\vec{\kappa}, j} \left\{ \sqrt{\hbar/8\pi A \omega_{sj}} [a_{sj}(\vec{\kappa}) + a_{sj}^{\dagger}(-\vec{\kappa})] \vec{\pi}_{sj}(\vec{r}) \right. \\ & \left. + \sqrt{\hbar/8\pi A \omega_{aj}} [a_{aj}(\vec{\kappa}) + a_{aj}^{\dagger}(-\vec{\kappa})] \vec{\pi}_{aj}(\vec{r}) \right\} \end{aligned} \quad (14a)$$

$$\begin{aligned} \dot{\vec{\pi}}(\vec{r}) = & \sum_{\vec{\kappa}, m} i \sqrt{\hbar \omega_{L\nu}/8\pi A} [a_m(\vec{\kappa}) - a_m^{\dagger}(-\vec{\kappa})] \dot{\vec{\pi}}_m(\vec{r}) \\ & + \sum_{\vec{\kappa}, j} \left\{ i \sqrt{\hbar \omega_{sj}/8\pi A} [a_{sj}(\vec{\kappa}) - a_{sj}^{\dagger}(-\vec{\kappa})] \dot{\vec{\pi}}_{sj}(\vec{r}) \right. \\ & \left. + i \sqrt{\hbar \omega_{aj}/8\pi A} [a_{aj}(\vec{\kappa}) - a_{aj}^{\dagger}(-\vec{\kappa})] \dot{\vec{\pi}}_{aj}(\vec{r}) \right\} \end{aligned} \quad (14b)$$

In Equations (14), we have introduced the area A of the interface, and the creation (annihilation) operators  $a_m^{\dagger}(\vec{\kappa})$  ( $a_m(\vec{\kappa})$ ) for the confined modes and  $a_{sj, aj}^{\dagger}(\vec{\kappa})$  ( $a_{sj, aj}(\vec{\kappa})$ ) for the symmetric and antisymmetric interface modes of phonons. They obey the commutation relations

$$[a_{\alpha}(\vec{\kappa}), a_{\beta}^{\dagger}(\vec{\kappa}')] = \delta_{\alpha\beta} \delta(\vec{\kappa} - \vec{\kappa}') \quad (15a)$$

$$[a_{\alpha}^{\dagger}(\vec{\kappa}), a_{\beta}^{\dagger}(\vec{\kappa}')] = [a_{\alpha}(\vec{\kappa}), a_{\beta}(\vec{\kappa}')] = 0 \quad (15b)$$

Using these commutators as well as the orthonormality relation for the eigenvectors, it is straightforward to show that

$$H_{ph} = H_{LO} + H_{IN} \quad (16a)$$

$$H_{LO} = \sum_{\vec{\kappa}, m} \hbar\omega_{L\vec{\kappa}} [a_m^{\dagger}(\vec{\kappa}) a_m(\vec{\kappa}) + \frac{1}{2}] \quad (16b)$$

$$H_{IN} = \sum_{\vec{\kappa}, j} \left\{ \hbar\omega_{sj}(\vec{\kappa}) [a_{sj}^{\dagger}(\vec{\kappa}) a_{sj}(\vec{\kappa}) + \frac{1}{2}] + \hbar\omega_{aj}(\vec{\kappa}) [a_{aj}^{\dagger}(\vec{\kappa}) a_{aj}(\vec{\kappa}) + \frac{1}{2}] \right\} \quad (16c)$$

The interaction energy for an electron interacting with the polarization at a point  $\vec{r}$  is represented by the Hamiltonian

$$H_{e-ph} = -e\phi(\vec{r}) \quad (17)$$

Regarded as a quantum mechanical operator, the scalar field can be expressed, with the help of (1), (4) and (5), as

$$\begin{aligned} \phi(\vec{r}) = & \sum_{\vec{\kappa}, m} \sqrt{\hbar/8\pi A\omega_{L\vec{\kappa}}} [a_m(\vec{\kappa}) + a_m^{\dagger}(-\vec{\kappa})] \phi_m(\vec{r}) \\ & + \sum_{\vec{\kappa}, j} \left\{ \sqrt{\hbar/8\pi A\omega_{sj}(\vec{\kappa})} [a_{sj}(\vec{\kappa}) + a_{sj}^{\dagger}(-\vec{\kappa})] \phi_{sj}(\vec{r}) \right. \\ & \left. + \sqrt{\hbar/8\pi A\omega_{aj}(\vec{\kappa})} [a_{aj}(\vec{\kappa}) + a_{aj}^{\dagger}(-\vec{\kappa})] \phi_{aj}(\vec{r}) \right\} \quad (18) \end{aligned}$$

Combining (10a), (11) and (18), we find after some algebra the interaction Hamiltonian

$$H_{e-ph} = H_{e-LO} + H_{e-IN} \quad (19)$$

$$H_{e-LO} = - \sum_{\vec{\kappa}} e^{i\vec{\kappa} \cdot \vec{\rho}} \left\{ \sum_{m=1,3,\dots} B_m(\kappa) \cos\left(\frac{m\pi z}{d}\right) [\hat{a}_m(\vec{\kappa}) + \hat{a}_m^{\dagger}(-\vec{\kappa})] \right\}$$



$$+ \sum_{m=2,4,\dots} B_m(\kappa) \sin\left(\frac{m\pi z}{d}\right) [\hat{a}_m(\vec{\kappa}) + \hat{a}_m^\dagger(-\vec{\kappa})] \quad (20)$$

$$H_{e-IN} = - \sum_{\vec{\kappa}, j} e^{i\vec{\kappa} \cdot \vec{\rho} - \kappa(|z| - d/2)} \left\{ B_{sj}(\vec{\kappa}) [\hat{a}_{s,j}(\vec{\kappa}) + \hat{a}_{s,j}^\dagger(-\vec{\kappa})] \right. \\ \left. - \operatorname{sgn}(z) B_{aj}(\kappa) [\hat{a}_{a,j}(\vec{\kappa}) + \hat{a}_{a,j}^\dagger(-\vec{\kappa})] \right\} \quad \text{for } |z| > d/2 \quad (21a)$$

$$H_{e-IN} = - \sum_{\vec{\kappa}, j} e^{i\vec{\kappa} \cdot \vec{\rho}} \left\{ B_{sj}(\kappa) \frac{\cosh(\kappa z)}{\cosh(\kappa d/2)} [\hat{a}_{s,j}(\vec{\kappa}) + \hat{a}_{s,j}^\dagger(-\vec{\kappa})] \right. \\ \left. - B_{aj}(\kappa) \frac{\sinh(\kappa z)}{\sinh(\kappa d/2)} [\hat{a}_{a,j}(\vec{\kappa}) + \hat{a}_{a,j}^\dagger(-\vec{\kappa})] \right\} \quad \text{for } |z| < d/2 \quad (21b)$$

The coefficients can be calculated from (10b) and (12). The results are

$$|B_m(\vec{\kappa})|^2 = \frac{e^2 \hbar}{8\pi A \omega_{Lv}} |C_m(\vec{\kappa})|^2 = \frac{4\pi e^2 \hbar \omega_{Lv}}{A d (\kappa^2 + \frac{m\pi}{d})} \left[ \frac{1}{\epsilon_{\omega\nu}} - \frac{1}{\epsilon_{0\nu}} \right] \quad (22a)$$

$$|B_{sj,aj}(\vec{\kappa})|^2 = \frac{e^2 \hbar}{8\pi A \omega_{sj,aj}(\vec{\kappa})} e^{-\kappa d} |C_{sj,aj}(\vec{\kappa})|^2, \quad (22b)$$

which leads to

$$|B_{sj}(\vec{\kappa})|^2 = \frac{\pi e^2}{A \kappa} - \hbar \omega_{sj}(\vec{\kappa}) / [\bar{\epsilon}_1 \tanh(\frac{\kappa d}{2}) + \bar{\epsilon}_2] \quad (23a)$$

$$|B_{aj}(\vec{\kappa})|^2 = \frac{\pi e^2}{A \kappa} - \hbar \omega_{aj}(\vec{\kappa}) / [\bar{\epsilon}_1 \coth(\frac{\kappa d}{2}) + \bar{\epsilon}_2] \quad (23b)$$

For convenience, we have defined in (23) the dielectric functions  $\bar{\epsilon}_\nu(\omega)$  by

$$\frac{1}{\bar{\epsilon}_\nu(\omega)} = \frac{1}{\epsilon_\nu(\omega) - \epsilon_{0\nu}} - \frac{1}{\epsilon_\nu(\omega) - \epsilon_{\omega\nu}} \quad (24)$$

#### 4. POLARON IN A DHS

We now proceed to calculate the polaron self-energy by perturbation theory. The unperturbed ground-state wave function is taken to be

$$|\psi_0\rangle = |\vec{k}, 0; n_{\vec{\kappa}, \beta}\rangle = \frac{1}{\sqrt{A}} e^{i\vec{k} \cdot \vec{\rho}} f(z) |n_{\vec{\kappa}, \beta}\rangle, \quad (25)$$

where electron states are characterized by a plane wave  $\vec{k}$  in the xy-plane and the ground state 0 in the z-direction. The electron wave vector is defined as  $\vec{k}_e = (\vec{k}, k_z)$  and the electron ground state wave function in a quantum well is given by

$$f(z) = \begin{cases} C \cos(k_z d/2) \exp[ik'_z(|z| - d/2)] & , \quad |z| \geq d/2 \\ C \cos(k_z z) & , \quad |z| < d/2 \end{cases} \quad (26)$$

The wave numbers  $k_z$  and  $k'_z$  are related to the electron subband energy  $E_\ell$  by

$$k_z = \sqrt{2m_e E_\ell / \hbar^2}, \quad k'_z = \sqrt{2m_e (V_0 - E_\ell) / \hbar^2} \quad (27)$$

with  $E_\ell$  determined by the transcendental equation

$$E_\ell = V_0 \cos^2\left(\frac{d}{2} \sqrt{2m_e E_\ell / \hbar^2}\right), \quad \ell = 1, 2, \dots \quad (28)$$

The normalization constant  $C$  is given by

$$C = \sqrt{2k'_z / (k'_z d + 2)} \quad (29)$$

The phonon state  $|n_{\vec{\kappa}, \beta}\rangle$  is specified by  $n$  phonons with wave vector  $\vec{k} = (\vec{\kappa}, q)$  and index  $\beta$  labeling the phonon mode, which may be one of the interface or confined LO modes.

The unperturbed energy of the electron in the ground state (25) can be written as

$$\epsilon_{\vec{k}_e}^{(0)} = \frac{\hbar k^2}{2m_e} + E_\ell + \sum_{\vec{\kappa}, \beta} (n + \frac{1}{2}) \hbar \omega_{\vec{\kappa}, \beta} \quad (30)$$

In the weak-coupling approximation, the corrections due to the electron-phonon interaction can be calculated by standard perturbation theory as

$$\begin{aligned} \epsilon_{\vec{k}_e} - \epsilon_{\vec{k}_e}^{(0)} &= \sum_{\vec{\kappa}, \beta} | \langle \vec{k}, 1_{\vec{\kappa}, \beta} | H_{e-ph} | \vec{k} - \vec{\kappa}, 0_{\vec{\kappa}, \beta} \rangle |^2 / \left[ \epsilon_{\vec{k}}^{(0)} - \epsilon_{\vec{k} - \vec{\kappa}}^{(0)} - \hbar \omega_{\vec{\kappa}, \beta} \right] \\ &= \Delta E_1 + \Delta E_2 + \Delta E_3, \end{aligned} \quad (31)$$

where we have assumed that transitions to or from a state with more than one phonon are negligible. Because we are only interested in the electron self-energy due to electron-phonon interactions, matrix elements corresponding to interband transitions are all ignored. With the Hamiltonian operators given by (19)-(21), and the wave function by (25), we find, after integration over the phonon momentum, that

$$\Delta E_1 = -\alpha_{L1} \hbar \omega_{L1} - \gamma_{L1} \hbar^2 k^2 / 2m_e \quad (32a)$$

$$\Delta E_2 = -\alpha_{L2} \hbar \omega_{L2} - \gamma_{L2} \hbar^2 k^2 / 2m_e \quad (32b)$$

$$\Delta E_3 = - \sum_j \int_0^\infty d\kappa \alpha_{IN}(\kappa) \hbar \omega_{sj}(\kappa) - \gamma_{IN} \hbar^2 k^2 / 2m_e \quad (32c)$$

The parameters  $\alpha$  and  $\gamma$  are related to the Fröhlich-type coupling constants involving the particular phonon modes indicated by the subscripts. Explicitly, the  $\alpha$ 's are given by

$$\alpha_{L1} = \frac{4\alpha_{F1}}{K_{p1}d} |C|^4 \sum_{m=1,3,\dots} \left(\frac{d}{m\pi}\right)^2 \left[1 + \frac{\cos(k_z d)}{1 - \left(\frac{2k_z d}{\mu\pi}\right)^2}\right] \left[\left(\frac{\pi}{K_{p1}d}\right)^2 - 1\right]^{-1} \ln \left|\frac{m\pi}{K_{p1}d}\right| \quad (33a)$$

$$\alpha_{L2} = \frac{4\alpha_{F2}}{\pi K_{p2}^2} |C|^4 \cos^4(k_z d/2) \int_0^\infty dt \frac{t^2 \ln|t|}{[t^2 + 4(k'_z/K_{p2})^2]^2 (t^2 - 1)} \quad (33b)$$

$$\alpha_{IN} = \alpha_{Fsj}(\kappa) |C|^4 / K_{psj}(\kappa) Q^2(\kappa) \left[\left(\frac{\kappa}{K_{psj}(\kappa)}\right)^2 + 1\right] \quad (33c)$$

and we have defined the function

$$Q^{-1}(\kappa) = \frac{2\cos^2(k_z d/2)}{\kappa + 2k'_z} + \frac{\tanh(\kappa d/2)}{\kappa} + \frac{2k_z \sin(k_z d)}{\kappa^2 + 4k_z^2} + \frac{\kappa}{\kappa^2 + 4k_z^2} \tanh\left(\frac{\kappa d}{2}\right) \cos(k_z d) \quad (34)$$

$K_{p1}^{-1}$  and  $K_{p2}^{-1}$  measure the polaron size due to the LO phonon in materials 1 and 2, respectively, while  $\alpha_{F1}$  ( $\alpha_{F2}$ ) denotes the usual LO-phonon-electron coupling constant of the Fröhlich type in material 1 (2). They are given by

$$\gamma_{L1} = \frac{4\alpha_{F1}}{K_{p1}d} |C|^4 \sum_{m=1,3,\dots} \left(\frac{d}{m\pi}\right)^2 \left[1 + \cos(k_z d) / \left[1 - \frac{2k_z d}{m\pi}\right]\right]^2$$

$$\times \int_0^\infty dt \, t / \left[t + \left(\frac{m\pi}{K_{p1}d}\right)^2\right] (t+1)^3 \quad (35a)$$

$$\gamma_{L2} = \frac{\alpha_{F2}}{k'_z K_{p2}} \cos^4(k_z d/2) \int_0^\infty dt \, t / (\sqrt{t} + 2k'_z/K_{p2})^2 (t+1)^3 \quad (35b)$$

$$\gamma_{IN} = \sum_j \int_0^\infty d\kappa \, 2\kappa^2 \alpha_{Fsj}(\kappa) |C|^4 / K_{psj}^3 Q^2(\kappa) [(\kappa/K_{psj})^2 + 1]^3 \quad (35c)$$

Substituting Eqs. (32) into (31) we find

$$\epsilon_{\vec{k}_e} - \epsilon_{\vec{k}_e}^{(0)} = -\alpha_{L1} \hbar\omega_{L1} - \alpha_{L2} \hbar\omega_{L2} - \sum_j \int_0^\infty d\kappa \, \alpha_{IN}(\kappa) \hbar\omega_{sj}(\kappa)$$

$$- \frac{\hbar^2 k^2}{2m_e} (\gamma_{L1} + \gamma_{L2} + \gamma_{IN}) \quad (36)$$

Combining (36) and the unperturbed energy (30), we obtain immediately the ground-state energy of the interacting electron-phonon system as

$$\epsilon_{\vec{k}_e} = \frac{\hbar^2 k^2}{2m_e^*} + E_g + \sum_{\vec{\kappa}, \beta} \left(n + \frac{1}{2}\right) \hbar\omega_{\kappa, \beta} - \Delta E \quad (37)$$

where

$$m_e^* = m_e / (1 - \gamma_{L1} - \gamma_{L2} - \gamma_{IN}) \quad (38)$$

is the polaron effective mass and

$$\Delta E = \alpha_{L1} \hbar\omega_{L1} + \alpha_{L2} \hbar\omega_{L2} + \sum_j \int_0^\infty d\kappa \, \alpha_{IN}(\kappa) \hbar\omega_{sj}(\kappa) \quad (39)$$

Equation (39) is now employed to compute the binding energy of a polaron in the GaAs quantum well of a GaAs/AlAs double heterostructure. The results are plotted as a function of the well width  $d$  in Fig. 2 in which contributions from the confined LO modes and interface modes are shown by dotted and dashed lines, respectively.




Figure 2. Polaron binding energy as a function of the well width. The dotted and dashed lines represent the contributions from confined LO and interface modes, respectively.

It is observed that for small  $d$ , our results are qualitatively different from what can be found in the literature. In fact, this is the first calculation that gives in the limit of  $d = 0$  the correct self-energy, the corresponding value in material 2. This is because we have included the contribution from confined LO phonons in material 2 in our Hamiltonian. The interface modes play an important role in the small- $d$  regime. Their influence quickly diminishes as  $d$  increases. It is important to note that interface phonons are responsible for the pinning effect observed in a quantum well on the transition energy of the polaron bound to a hydrogenic impurity.<sup>11</sup> This is discussed in the following section.

##### 5. TRANSITION ENERGY OF A BOUND MAGNETOPOLARON

Recent experiments<sup>11,12</sup> reveal that the resonant effect of the electron-phonon-interaction on the  $1s-2p$  transition energy of a magnetopolaron bound to a hydrogenic impurity in GaAs/AlGaAs multiple-quantum-well structures exhibits the pinning phenomenon at an energy below the TO-phonon energy. To our knowledge, the origin of this surprising effect has not been understood.<sup>13</sup> Here we attempt to calculate this transition energy in the GaAs well of a GaAs/AlGaAs DHS.

Consider a donor impurity at the center of the well as in the case of the sample used in the experiments. A magnetic field is applied along the axis of the quantum well. The total Hamiltonian can be written as

$$H = H_e + H_{ph} + H_{e-ph} \quad (40)$$

where

$$H_e = [\vec{p} - \frac{e}{c} \vec{A}]^2 / 2m_e - \frac{e^2}{\epsilon_0 r} + V_B(z) \quad (41a)$$

represents the energy of a hydrogenic impurity confined in the square well given by

$$V_B(z) = \begin{cases} 0 & , \quad |z| \leq d/2 \\ V_0 & , \quad |z| > d/2 \end{cases} \quad (41b)$$

the free phonon energy  $H_{ph}$  is given by (16) and the interaction  $H_{e-ph}$  by (19)-(21).

The unperturbed energy levels of the hydrogenic impurity described by (41a) are calculated variationally. For the trial wave function, we take

$$\psi = f(z) G(\rho, z, \phi) \quad (42)$$

The function  $f(z)$  is the solution to the square-well problem and is given by

$$f(z) = \begin{cases} \cos(k_z z) & , \quad z \leq d/2 \\ Ae^{-k'_z z} & , \quad z > d/2 \end{cases} \quad (43)$$

where  $k_z$  and  $k'_z$  are defined in (27) with  $E_j$  representing the energy of the first subband. The function  $G(\rho, z, \phi)$  describes the internal states of the donor and is chosen to reflect the symmetry properties of the system,

$$G(\rho, z, \phi) = \rho^{|m|} e^{im\phi} e^{-\gamma\rho^2/4} e^{-\gamma\xi^2 z^2/4} \quad (44)$$

where  $\xi$  is the variational parameter to be determined by minimizing the ground-state energy.

The interaction  $H_{e-ph}$  is then treated as a perturbation, and the energy levels are calculated by means of Wigner-Brillouin perturbation theory. The result is<sup>14</sup>

$$E_i(B) = E_i(B) + \frac{1}{(2\pi)^2} \sum_n \int d^2\kappa \frac{|\langle n | H_{e-ph} | i \rangle|^2}{E_i(B) - E_n(B) - \hbar\omega(\vec{\kappa})} \quad (45)$$

where  $E_i(B)$  stands for the corresponding unperturbed energy. The calculation is rather complicated, and detailed discussion of the procedures will be published elsewhere.<sup>15</sup> Here we just mention that the Hilbert space has been truncated to only the lowest three states in our second-order perturbation term. In other words, only  $1s$ ,  $2p_+$  and  $2p_-$  are included. For the particular samples used in these

experiments,<sup>11,12</sup> all other states including  $2p_0$  are pushed up by the quantum well way too high because of the broken symmetry and cannot be reached with presently-available magnetic fields. Furthermore, interactions of the electron with lattice vibrations outside the well under consideration are ignored, since they are important only for a very narrow ( $d \leq 10$  Å) well, while  $d$  is larger than 100 Å in the experiments considered here.




Figure 3.  $1s-2p_+$  transition energy as a function of the magnetic field  $B$ . Data are taken at SUNY-Buffalo and at the Francis Bitter National Magnet Laboratory in a series of measurements as indicated at the lower right corner. The heavy solid line represent results from the present theory and the straight line is the result without electron-phonon interaction.

The  $1s-2p_+$  transition energy calculated by computer iteration is plotted as a function of the field  $B$  in Fig. 3 along with the newly-improved set of data.<sup>12</sup> It is remarkable that the theoretical curve breaks into three branches separated by two gaps, in good agreement with experiments. The occurrence of gaps reflects the energy level repelling caused by the strong resonance interaction when the transition energy matches the phonon energy. Since the electron does not couple to TO-phonons, this pinning phenomenon must be due to the electron interaction with interface phonons which oscillate at frequencies between the bulk  $\omega_L$  and  $\omega_T$ . The

interface-phonon coupling alone, however, is not adequate for a complete understanding of this phenomenon. There are still small discrepancies between experimental data and the theory, especially in the lower branch near the gap. Minor corrections such as the nonparabolic effect on the effective mass can probably account for the discrepancies. This and other possible corrections will be discussed in a forthcoming article.<sup>15</sup>

## 6. ACKNOWLEDGMENTS

We are grateful to B. D. McCombe and J. P. Cheng for providing us with experimental data prior to their publication. This research was supported in part by the U. S. Office of Naval Research.

## 7. REFERENCES

1. R. Lassnig, Phys. Rev. B 30, 7132 (1984).
2. L. Wendler, Phys. Stat. Solidi B 129, 513 (1985).
3. R. Fuchs and K. L. Kliewer, Phys. Rev. 140, A2076 (1965).
4. J. J. Licari and Evrard, Phys. Rev. B 15, 2254 (1977).
5. X. X. Liang, S. W. Gu and D. L. Lin, Phys. Rev. B 34, 1251 (1986).
6. C. Y. Chen, T. Z. Ding and D. L. Lin, Phys. Rev. B 35, 4398 (1987); C. Y. Chen, D. L. Lin and T. Z. Ding, Phys. Rev. B 36, 9816 (1987).
7. S. W. Gu, X. J. Kong and C. W. Wei, Phys. Rev. B 36, 7977 (1987).
8. R. Chen, D. L. Lin and T. F. George, Phys. Rev. B 41, 1435 (1990); D. L. Lin, R. Chen and T. F. George, Solid State Commun. (in press).
9. N. Mori and T. Ando, Phys. Rev. 40, 6175 (1989).
10. J. E. Zucker, A. Pinczuk, D. S. Chemla, A. Gossard and W. Wiegmann, Phys. Rev. Lett. 53, 1280 (1984).
11. Y. H. Change, B. D. McCombe, J. M. Mercy, A. A. Reeder, J. Ralston and G. A. Wicks, Phys. Rev. Lett. 61, 1408 (1988).
12. J. P. Cheng, B. D. McCombe and G. Brozak, to be published.
13. J. P. Cheng and B. D. McCombe, Phys. Rev. Lett. 62, 1925 (1989).
14. D. R. Cohn, D. M. Larsen and B. Lax, Phys. Rev. B 6, 1367 (1972).
15. D. L. Lin, R. Chen and T. F. George, to be published.



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSROC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
		Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen  
Hughes Research Laboratory  
3011 Malibu Canyon Road  
Malibu, California 90265

Dr. J. H. Weaver  
Department of Chemical Engineering  
and Materials Science  
University of Minnesota  
Minneapolis, Minnesota 55455

Dr. A. Reisman  
Microelectronics Center of North Carolina  
Research Triangle Park, North Carolina  
27709

Dr. M. Grunze  
Laboratory for Surface Science and  
Technology  
University of Maine  
Orono, Maine 04469

Dr. J. Butler  
Naval Research Laboratory  
Code 6115  
Washington D.C. 20375-5000

Dr. L. Interante  
Chemistry Department  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. Irvin Heard  
Chemistry and Physics Department  
Lincoln University  
Lincoln University, Pennsylvania 19352

Dr. K.J. Klaubunde  
Department of Chemistry  
Kansas State University  
Manhattan, Kansas 66506

Dr. C. B. Harris  
Department of Chemistry  
University of California  
Berkeley, California 94720

Dr. F. Kutzler  
Department of Chemistry  
Box 5055  
Tennessee Technological University  
Cookeville, Tennessee 38501

Dr. D. DiLella  
Chemistry Department  
George Washington University  
Washington D.C. 20052

Dr. R. Reeves  
Chemistry Department  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. Steven M. George  
Stanford University  
Department of Chemistry  
Stanford, CA 94305

Dr. Mark Johnson  
Yale University  
Department of Chemistry  
New Haven, CT 06511-8118

Dr. W. Knauer  
Hughes Research Laboratory  
3011 Malibu Canyon Road  
Malibu, California 90265

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. G. A. Somorjai  
Department of Chemistry  
University of California  
Berkeley, California 94720

Dr. J. Murday  
Naval Research Laboratory  
Code 6170  
Washington, D.C. 20375-5000

Dr. J. B. Hudson  
Materials Division  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. Theodore E. Madey  
Surface Chemistry Section  
Department of Commerce  
National Bureau of Standards  
Washington, D.C. 20234

Dr. J. E. Demuth  
IBM Corporation  
Thomas J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, New York 10598

Dr. M. G. Lagally  
Department of Metallurgical  
and Mining Engineering  
University of Wisconsin  
Madison, Wisconsin 53706

Dr. R. P. Van Duyne  
Chemistry Department  
Northwestern University  
Evanston, Illinois 60637

Dr. J. M. White  
Department of Chemistry  
University of Texas  
Austin, Texas 78712

Dr. D. E. Harrison  
Department of Physics  
Naval Postgraduate School  
Monterey, California 93940

Dr. R. L. Park  
Director, Center of Materials  
Research  
University of Maryland  
College Park, Maryland 20742

Dr. W. T. Peria  
Electrical Engineering Department  
University of Minnesota  
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson  
Department of Metallurgy and  
Materials Science  
Massachusetts Institute of Technology  
Cambridge, Massachusetts 02139

Dr. S. Sibener  
Department of Chemistry  
James Franck Institute  
5640 Ellis Avenue  
Chicago, Illinois 60637

Dr. Arnold Green  
Quantum Surface Dynamics Branch  
Code 3817  
Naval Weapons Center  
China Lake, California 93555

Dr. A. Wold  
Department of Chemistry  
Brown University  
Providence, Rhode Island 02912

Dr. S. L. Bernasek  
Department of Chemistry  
Princeton University  
Princeton, New Jersey 08544

Dr. W. Kohn  
Department of Physics  
University of California, San Diego  
La Jolla, California 92037

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter  
Code 6170  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. Richard Colton  
Code 6170  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. Dan Pierce  
National Bureau of Standards  
Optical Physics Division  
Washington, D.C. 20234

Dr. R. Stanley Williams  
Department of Chemistry  
University of California  
Los Angeles, California 90024

Dr. R. P. Messmer  
Materials Characterization Lab.  
General Electric Company  
Schenectady, New York 22217

Dr. Robert Gomer  
Department of Chemistry  
James Franck Institute  
5640 Ellis Avenue  
Chicago, Illinois 60637

Dr. Ronald Lee  
R301  
Naval Surface Weapons Center  
White Oak  
Silver Spring, Maryland 20910

Dr. Paul Schoen  
Code 6190  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. John T. Yates  
Department of Chemistry  
University of Pittsburgh  
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene  
Code 520  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. L. Kesmodel  
Department of Physics  
Indiana University  
Bloomington, Indiana 47403

Dr. K. C. Janda  
University of Pittsburgh  
Chemistry Building  
Pittsburgh, PA 15260

Dr. E. A. Irene  
Department of Chemistry  
University of North Carolina  
Chapel Hill, North Carolina 27514

Dr. Adam Heller  
Bell Laboratories  
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann  
Department of Chemistry  
University of Southampton  
Southampton SO9 5NH  
UNITED KINGDOM

Dr. H. Tachikawa  
Chemistry Department  
Jackson State University  
Jackson, Mississippi 39217

Dr. John W. Wilkins  
Cornell University  
Laboratory of Atomic and  
Solid State Physics  
Ithaca, New York 14853

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis  
Department of Physics  
University of California  
Irvine, California 92664

Dr. D. Ramaker  
Chemistry Department  
George Washington University  
Washington, D.C. 20052

Dr. J. C. Hemminger  
Chemistry Department  
University of California  
Irvine, California 92717

Dr. T. F. George  
Chemistry Department  
University of Rochester  
Rochester, New York 14627

Dr. G. Rubloff  
IBM  
Thomas J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, New York 10598

Dr. Horia Metiu  
Chemistry Department  
University of California  
Santa Barbara, California 93106

Dr. W. Goddard  
Department of Chemistry and Chemical  
Engineering  
California Institute of Technology  
Pasadena, California 91125

Dr. P. Hansma  
Department of Physics  
University of California  
Santa Barbara, California 93106

Dr. J. Baldeschwieler  
Department of Chemistry and  
Chemical Engineering  
California Institute of Technology  
Pasadena, California 91125

Dr. J. T. Keiser  
Department of Chemistry  
University of Richmond  
Richmond, Virginia 23173

Dr. R. W. Plummer  
Department of Physics  
University of Pennsylvania  
Philadelphia, Pennsylvania 19104

Dr. E. Yeager  
Department of Chemistry  
Case Western Reserve University  
Cleveland, Ohio 44106

Dr. N. Winograd  
Department of Chemistry  
Pennsylvania State University  
University Park, Pennsylvania 16802

Dr. Roald Hoffmann  
Department of Chemistry  
Cornell University  
Ithaca, New York 14853

Dr. A. Steckl  
Department of Electrical and  
Systems Engineering  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. G.H. Morrison  
Department of Chemistry  
Cornell University  
Ithaca, New York 14853